

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING
FOR ANIMAL TREATMENT ONLY

VALBAZEN[®]
BROAD SPECTRUM SHEEP LAMB AND GOAT DRENCH

ACTIVE CONSTITUENT:
ALBENDAZOLE 19 mg/mL



For the control of susceptible gastrointestinal roundworms (except whipworms), large lungworms, tapeworm and to aid in the control of adult liver fluke.

Also to reduce the output of viable worm and fluke eggs.

5 L

F-LA510489/00

COOPERS[®]
EST. 1843



VALBAZEN®

BROAD SPECTRUM SHEEP LAMB AND GOAT DRENCH

For the control of worms susceptible to benzimidazole type drenches including worms resistant to levamisole/morantel and organophosphate type drenches.

Roundworms (mature and immature)

Barbers pole worm
Stomach hair worm
Small brown stomach worm
Black scour worm
Small intestinal worm
Thin-necked intestinal worm
Hookworm
Nodule worm
Large bowel worm
Large mouthed bowel worm

Lungworms: Large lungworm

Tapeworms: Heads and Segments

Liver Fluke:

Aids in control of adult liver fluke.

Ovicidal:

Reduces the output of viable worm and fluke eggs.

Internal parasites can develop resistance to drenches. This drench is not effective against worms resistant to benzimidazole type drenches. Ask your local veterinary advisor for recommended parasite management programs to reduce resistance development. A drench resistance test should be conducted before any drench is used.

DIRECTIONS FOR USE:

Restraints:

DO NOT USE in animals which are producing or may in the future produce milk or milk products for human consumption.

Precautions:

Shake container before use.

Management practices should ensure that all drenching procedures are correctly carried out and bodyweights accurately assessed.

During the early part of pregnancy all medication should be administered with care and it is particularly important not to exceed the recommended dose at this time.

DOSE RATES - SHEEP, LAMBS, GOATS

Roundworms - Lungworms - Tapeworms: 1 mL/5 kg bodyweight

Roundworms - Lungworms - Tapeworms

- Adult Liver Fluke: 1 mL/4 kg bodyweight

Bodyweight	Roundworms Lungworms Tapeworms	5 L Container Treats	Roundworms Lungworms Tapeworms Adult Liver Fluke	5 L Container Treats
10 kg	2 mL	2,500	2.5 mL	2,000
15 kg	3 mL	1,666	4.0 mL	1,250
20 kg	4 mL	1,250	5.0 mL	1,000
25 kg	5 mL	1,000	6.5 mL	769
30 kg	6 mL	833	7.5 mL	666
35 kg	7 mL	714	9.0 mL	555
40 kg	8 mL	625	10.0 mL	500
45 kg	9 mL	555	11.5 mL	434
50 kg	10 mL	500	12.5 mL	400
55 kg	11 mL	454	14.0 mL	357
60 kg	12 mL	416	15.0 mL	333
65 kg	13 mL	384	16.5 mL	303
70 kg	14 mL	357	17.5 mL	285
75 kg	15 mL	333	19.0 mL	263

Animals over 75 kg to be dosed at rates above. Dose to heaviest animal in each group (ewes/does, rams/bucks, lambs/kids), do not underdose. Animals should be weighed.

VALBAZEN AIDS IN THE CONTROL OF ADULT LIVER FLUKE

VALBAZEN reduces the output of viable fluke eggs. Contamination of pastures will be reduced by avoiding grazing recognised fluke areas for 24 hours after dosing. A second treatment with VALBAZEN one month after the first will assist to prevent contamination of pastures and will minimise chances of reinfestation. Avoid placing treated stock on clean pastures for a minimum of 4 hours after dosing.

CLEAN DRENCHING EQUIPMENT BEFORE AND AFTER USE.

IMPORTANT NEW INFORMATION

CSIRO trials have shown increased worm killing activity by Valbazen when feed is withdrawn from sheep prior to and after dosing. Better results can be achieved when using Valbazen if sheep are held in holding areas or yards with no feed but with free access to water for 24 hours before dosing and 8 hours after. Restricting feed should not be practised on sheep that are heavily pregnant, severely stressed or in poor condition.

WITHHOLDING PERIODS

MEAT: DO NOT USE less than 10 days before slaughter for human consumption.

MILK: DO NOT USE in animals which are producing or may in the future produce milk or milk products for human consumption.

TRADE ADVICE:

EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 10 days before slaughter for export. The ESI on this label was correct at the time of label approval. Before using this product, confirm the current ESI from the manufacturer on 1800 226 511 or the APVMA website (www.apvma.gov.au/residues/ESI.shtml).

For further information contact Coopers Customer Service on 1 800 226 511.

SAFETY DIRECTIONS

Harmful if absorbed by skin contact or swallowed. Avoid contact with skin. Wash hands after use.

Additional information is in the Material Safety Data Sheet.

SPECIALIST ADVICE IN EMERGENCY ONLY

Coopers Animal Health (Intervet Australia Pty Ltd)

1800 226 511

ALL HOURS - AUSTRALIA WIDE

Warranty

Intervet Australia Pty Ltd (IAPL), also known as MSD Animal Health, warrants that this product is of merchantable quality and fit for its intended purpose. IAPL's liability for any loss, including consequential losses or injury caused by any act or omission, including negligent acts or omissions, by IAPL or its agent is limited to replacing or repairing the product at the option of IAPL. If possible, a sample of any product causing concern should be retained or delivered to IAPL within 30 days for a scientific examination.

DISPOSAL

This container can be recycled if it is clean, dry, free of visible residues and has the **drumMUSTER** logo visible. Triple or pressure rinse container for disposal. Dispose of rinsate or any undiluted chemical according to State legislative requirements. Wash outside of the container and the cap.

Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any **drumMUSTER** collection or similar container management program site. The cap should not be replaced but may be taken separately.

STORAGE: Store Below 30°C (Room Temperature).

SEE BELOW FOR BATCH AND EXPIRY



COOPERS ANIMAL HEALTH

(Intervet Australia Pty Ltd)

91-105 Harpin Street

Bendigo East, VIC 3550

APVMA Approval No. 45788/53022

® Registered Trademark

® Yellow (PANTONE® 123) the predominant colour of Coopers packaging is a Registered Trademark.

PANTONE is a registered trademark of Pantone, Inc.

© 2012



BUS10489/00