

observed in the present study is due to inositol depletion or to the avoidance of allergenic foods, the results suggest that some patients with bipolar disorder respond to dietary modifications.

Shaldubina A, et al. Inositol deficiency diet and lithium effects. *Bipolar Disord.* 2006;8: 152-159.

High iodine intake associated with thyroiditis and hypothyroidism

Salt has been iodized in China since 1996, resulting in an increase in iodine intake throughout the country. In a 1999 study, researchers observed an increase in the prevalence of autoimmune thyroiditis, overt hypothyroidism, and subclinical hypothyroidism with increasing iodine intake in cohorts from three regions of China with different levels of iodine intake: "mildly deficient" (median urinary iodine excretion, 84 mcg/L), "more than adequate" (median, 243 mcg/L), and "excessive" (median, 651 mcg/L). Of the 3,761 subjects enrolled in the original study, 3,018 (80.2%) participated in a five-year, follow-up study. During




the follow-up period, among subjects with mildly deficient iodine intake, more than adequate intake, and excessive intake, the cumulative incidence of autoimmune thyroiditis was 0.2%, 1.0%, and 1.3%, respectively; that of subclinical hypothyroidism, 0.2%, 2.6%, and 2.9%, respectively; and that of overt hypothyroidism, 0.2%, 0.5%, and 0.3%, respectively. The differences in incidence for mildly deficient vs. more than adequate or excessive intake were statistically significant for autoimmune thyroiditis ($p = 0.01$ to 0.03) and for subclinical hypothyroidism ($p < 0.001$). The authors concluded that more than adequate or excessive iodine intake may lead to autoimmune thyroiditis and hypothyroidism.

Comment: Iodine deficiency remains an important problem in some parts of the world, and iodine intake should be increased in people whose intake is inadequate. High-dose iodine therapy also has a role in clinical medicine, particularly in the treatment of fibrocystic breast disease. However, people taking large amounts of iodine should be monitored for the development of thyroid abnormalities.

Teng W, et al. Effect of iodine intake on thyroid diseases in China. *N Engl J Med.* 2006;354:2783-2793.

N-acetylcysteine for polycystic ovary syndrome

One hundred-fifty overweight or obese infertile women (mean age, 29 years; range, 18-39 years) with polycystic ovary syndrome (PCOS) who had failed to ovulate after treatment



magnetico Sleep Pads

... FOR PROVEN PAIN RELIEF from

FIBROMYALGIA & CHRONIC FATIGUE

STUDY #1

U of VA - Double Blind
111 Patients - Placebo Controlled

Magnetico Sleep Pad Users
✓ Showed Significant,
Lasting Improvement in
FUNCTION & PAIN REDUCTION

STUDY #2

Dr. G Lewis - Clinical, Double Blind
29 Patients - Placebo Controlled

Magnetico Sleep Pad Users
Showed 2.4X Improvement in:
✓ **PAIN REDUCTION**
✓ **IMPROVED SLEEP/ LESS FATIGUE**
✓ **ABILITY TO WORK LONGER HOURS**

NOTE: Dr Bonlie has since developed two stronger model Magnetico Sleep Pads as recommended by the U of VA upon evaluating the study results. Magnetico Sleep Pads are placed **BENEATH** your mattress... so any style will do! **"Six-Month Satisfaction Guarantee". CALL TODAY for your**

FREE 24 - page Booklet & Lecture DVD, 800-265-1119 www.magneticosleep.com

Copyright of Townsend Letter for Doctors & Patients is the property of Townsend Letter Group and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.